

A bilinear Airy-estimate with application to gKdV-3

Axel Grünrock
 Fachbereich Mathematik
 Bergische Universität - Gesamthochschule Wuppertal
 Gaußstraße 20
 D-42097 Wuppertal
 Germany
 e-mail Axel.Gruenrock@math.uni-wuppertal.de

Abstract

The Fourier restriction norm method is used to show local wellposedness for the Cauchy-Problem

$$u_t + u_{xxx} + (u^4)_x = 0, \quad u(0) = u_0 \in H_x^s(\mathbf{R}), \quad s > -\frac{1}{6}$$

for the generalized Korteweg-deVries equation of order three, for short gKdV-3. For real valued data $u_0 \in L_x^2(\mathbf{R})$ global wellposedness follows by the conservation of the L^2 -norm. The main new tool is a bilinear estimate for solutions of the Airy-equation.

The purpose of this note is to establish local wellposedness of the Cauchy-Problem

$$u_t + u_{xxx} + (u^4)_x = 0, \quad u(0) = u_0 \in H_x^s(\mathbf{R}), \quad s > -\frac{1}{6}$$

for the generalized Korteweg-deVries equation of order three, for short gKdV-3. So far, local wellposedness of this problem is known for data $u_0 \in H_x^s(\mathbf{R})$, $s \geq \frac{1}{12}$. This was shown by Kenig, Ponce and Vega in 1993, see Theorem 2.6 in [KPV93]. Here we extend this result to data $u_0 \in H_x^s(\mathbf{R})$, $s > -\frac{1}{6}$. A standard scaling argument suggests that this is optimal (up to the endpoint). For real valued data $u_0 \in L_x^2(\mathbf{R})$ we obtain global wellposedness by the conservation of the L^2 -norm.

By the Fourier restriction norm method introduced in [B93] and further developed in [KPV96] and [GTV97] matters reduce to the proof of the estimate

$$\|\partial_x \prod_{i=1}^4 u_i\|_{X_{s,b'}} \leq c \prod_{i=1}^4 \|u_i\|_{X_{s,b}} \quad (1)$$

for suitable values of s , b and b' . Here the space $X_{s,b}$ is the completion of the Schwartz class $\mathcal{S}(\mathbf{R}^2)$ with respect to the norm

$$\|u\|_{X_{s,b}} = \left(\int d\xi d\tau <\tau - \xi^3 >^{2b} <\xi >^{2s} |\mathcal{F}u(\xi, \tau)|^2 \right)^{\frac{1}{2}},$$

where $\mathcal{F}$ denotes the Fourier transform in both variables. The main new tool for the proof of (1) is the following bilinear Airy-estimate:

Lemma 1 Let I^s denote the Riesz potential of order $-s$ and let $I_-^s(f, g)$ be defined by its Fourier transform (in the space variable):

$$\mathcal{F}_x I_-^s(f, g)(\xi) := \int_{\xi_1 + \xi_2 = \xi} d\xi_1 |\xi_1 - \xi_2|^s \mathcal{F}_x f(\xi_1) \mathcal{F}_x g(\xi_2).$$

Then we have

$$\|I_-^{\frac{1}{2}} I_-^{\frac{1}{2}}(e^{-t\partial^3} u_1, e^{-t\partial^3} u_2)\|_{L_{xt}^2} \leq c \|u_1\|_{L_x^2} \|u_2\|_{L_x^2}.$$

Proof: We will write for short $\hat{u}$ instead of $\mathcal{F}_x u$ and $\int_* d\xi_1$ for $\int_{\xi_1 + \xi_2 = \xi} d\xi_1$. Then, using Fourier-Plancherel in the space variable we obtain:

$$\begin{aligned} & \|I_-^{\frac{1}{2}} I_-^{\frac{1}{2}}(e^{-t\partial^3} u_1, e^{-t\partial^3} u_2)\|_{L_{xt}^2}^2 \\ &= c \int d\xi |\xi| dt \left| \int_* d\xi_1 |\xi_1 - \xi_2|^{\frac{1}{2}} e^{it(\xi_1^3 + \xi_2^3)} \hat{u}_1(\xi_1) \hat{u}_2(\xi_2) \right|^2 \\ &= c \int d\xi |\xi| dt \int_* d\xi_1 d\eta_1 e^{it(\xi_1^3 + \xi_2^3 - \eta_1^3 - \eta_2^3)} (|\xi_1 - \xi_2| |\eta_1 - \eta_2|)^{\frac{1}{2}} \prod_{i=1}^2 \hat{u}_i(\xi_i) \overline{\hat{u}_i(\eta_i)} \\ &= c \int d\xi |\xi| \int_* d\xi_1 d\eta_1 \delta(\eta_1^3 + \eta_2^3 - \xi_1^3 - \xi_2^3) (|\xi_1 - \xi_2| |\eta_1 - \eta_2|)^{\frac{1}{2}} \prod_{i=1}^2 \hat{u}_i(\xi_i) \overline{\hat{u}_i(\eta_i)} \\ &= c \int d\xi |\xi| \int_* d\xi_1 d\eta_1 \delta(3\xi(\eta_1^2 - \xi_1^2 + \xi(\xi_1 - \eta_1))) (|\xi_1 - \xi_2| |\eta_1 - \eta_2|)^{\frac{1}{2}} \prod_{i=1}^2 \hat{u}_i(\xi_i) \overline{\hat{u}_i(\eta_i)}. \end{aligned}$$

Now we use $\delta(g(x)) = \sum_n \frac{1}{|g'(x_n)|} \delta(x - x_n)$, where the sum is taken over all simple zeros of g , in our case:

$$g(x) = 3\xi(x^2 + \xi(\xi_1 - x) - \xi_1^2)$$

with the zeros $x_1 = \xi_1$ and $x_2 = \xi - \xi_1$, hence $g'(x_1) = 3\xi(2\xi_1 - \xi)$ respectively $g'(x_2) = 3\xi(\xi - 2\xi_1)$. So the last expression is equal to

$$\begin{aligned} & c \int d\xi |\xi| \int_* d\xi_1 d\eta_1 \frac{1}{|\xi| |2\xi_1 - \xi|} \delta(\eta_1 - \xi_1) (|\xi_1 - \xi_2| |\eta_1 - \eta_2|)^{\frac{1}{2}} \prod_{i=1}^2 \hat{u}_i(\xi_i) \overline{\hat{u}_i(\eta_i)} \\ &+ c \int d\xi |\xi| \int_* d\xi_1 d\eta_1 \frac{1}{|\xi| |2\xi_1 - \xi|} \delta(\eta_1 - (\xi - \xi_1)) (|\xi_1 - \xi_2| |\eta_1 - \eta_2|)^{\frac{1}{2}} \prod_{i=1}^2 \hat{u}_i(\xi_i) \overline{\hat{u}_i(\eta_i)} \\ &= c \int d\xi \int_* d\xi_1 \prod_{i=1}^2 |\hat{u}_i(\xi_i)|^2 + c \int d\xi \int_* d\xi_1 \hat{u}_1(\xi_1) \overline{\hat{u}_1(\xi_2)} \hat{u}_2(\xi_2) \overline{\hat{u}_2(\xi_1)} \\ &\leq c \left(\prod_{i=1}^2 \|u_i\|_{L_x^2}^2 + \|\hat{u}_1 \hat{u}_2\|_{L_\xi^1}^2 \right) \leq c \prod_{i=1}^2 \|u_i\|_{L_x^2}^2. \end{aligned}$$

□

Arguing as in the proof of Lemma 2.3 in [GTV97] we get the following

Corollary 1 Let $b > \frac{1}{2}$. Then the following estimate holds true:

$$\|I_-^{\frac{1}{2}} I_-^{\frac{1}{2}}(u, v)\|_{L_{xt}^2} \leq c \|u\|_{X_{0,b}} \|v\|_{X_{0,b}}$$

In the next Lemma, some well known Strichartz type estimates for the Airy equation are gathered in terms of $X_{s,b}$ -norms:

Lemma 2 For $b > \frac{1}{2}$ the following estimates are valid:

$$i) \|u\|_{L_t^p(\dot{H}_x^{s,q})} \leq c\|u\|_{X_{0,b}}, \text{ whenever } 0 \leq s = \frac{1}{p} \leq \frac{1}{4} \text{ and } \frac{1}{q} = \frac{1}{2} - \frac{2}{p},$$

$$ii) \|u\|_{L_t^p(L_x^q)} \leq c\|u\|_{X_{0,b}}, \text{ whenever } 0 < \frac{1}{q} = \frac{1}{2} - \frac{3}{p} \leq \frac{1}{2}.$$

Quotation/proof: Theorem 2.1 in [KPV91] gives in the case of the Airy-equation

$$\|e^{-t\partial^3} u_0\|_{L_t^p(\dot{H}_x^{s,q})} \leq c\|u_0\|_{L_x^2},$$

provided $0 \leq s = \frac{1}{p} \leq \frac{1}{4}$ and $\frac{1}{q} = \frac{1}{2} - \frac{2}{p}$. Now Lemma 2.3 in [GTV97] is applied to obtain

$$\|u\|_{L_t^p(\dot{H}_x^{s,q})} \leq c\|u\|_{X_{0,b}}, \quad b > \frac{1}{2} \quad (2)$$

for the same values of s , p and q . From this ii) follows by Sobolev's embedding theorem (in the space variable). Especially we have

$$\|u\|_{L_{xt}^8} \leq c\|u\|_{X_{0,b}}, \quad b > \frac{1}{2},$$

which, interpolated with the trivial case, gives

$$\|u\|_{L_{xt}^4} \leq c\|u\|_{X_{0,b}}, \quad b > \frac{1}{3}.$$

Now let us see how to replace $\dot{H}_x^{s,q}$ by $H_x^{s,q}$ in (2) in the endpoint case, i. e. $s = \frac{1}{p} = \frac{1}{4}$, $q = \infty$: Using the projections $p = \mathcal{F}_x^{-1} \chi_{\{|\xi| \leq 1\}} \mathcal{F}_x$ and $P = Id - p$ we have

$$\|u\|_{L_t^4(H_x^{\frac{1}{4},\infty})} \leq \|Pu\|_{L_t^4(H_x^{\frac{1}{4},\infty})} + \|pu\|_{L_t^4(H_x^{\frac{1}{4},\infty})} =: I + II.$$

For I we use (2) to obtain

$$I \leq c\|I^{-\frac{1}{4}} J^{\frac{1}{4}} Pu\|_{X_{0,b}} \leq c\|u\|_{X_{0,b}},$$

while for II by Sobolev's embedding theorem we get

$$II \leq c\|pu\|_{L_t^4(H_x^{\frac{1}{2}+,4})} \leq c\|pu\|_{X_{\frac{1}{2}+,b}} \leq c\|u\|_{X_{0,b}}.$$

This gives i) in the endpoint case, from which the general case follows by interpolation with Sobolev's embedding theorem (in the time variable). $\square$

Remark: The endpoint case in ii) is also valid - see e. g. Lemma 3.29 in [KPV93] - but we shall not make use of this here.

Now we are prepared to prove the crucial nonlinear estimate:

Theorem 1 For $0 \geq s > -\frac{1}{6}$, $-\frac{1}{2} < b' < s - \frac{1}{3}$ and $b > \frac{1}{2}$ the estimate (1) is valid.

Proof: Writing $f_i(\xi, \tau) = \langle \tau - \xi^3 \rangle^b \langle \xi \rangle^s \mathcal{F}u_i(\xi, \tau)$, $1 \leq i \leq 4$, we have

$$\|\partial_x \prod_{i=1}^4 u_i\|_{X_{s,b'}} = c\|\langle \tau - \xi^3 \rangle^{b'} \langle \xi \rangle^s |\xi| \int d\nu \prod_{i=1}^4 \langle \tau_i - \xi_i^3 \rangle^{-b} \langle \xi_i \rangle^{-s} f_i(\xi_i, \tau_i)\|_{L_{\xi,\tau}^2},$$

where $d\nu = d\xi_1 \dots d\xi_3 d\tau_1 \dots d\tau_3$ and $\sum_{i=1}^4 (\xi_i, \tau_i) = (\xi, \tau)$. Now the domain of integration is divided into the regions A , B and $C = (A \cup B)^c$, where in A we assume $|\xi_{max}| \leq c$. (Here ξ_{max} is defined by $|\xi_{max}| = \max_{i=1}^4 |\xi_i|$, similarly ξ_{min} .) Then for the region A we have the upper bound

$$\begin{aligned} & c\|\int d\nu \prod_{i=1}^4 \langle \tau_i - \xi_i^3 \rangle^{-b} f_i(\xi_i, \tau_i)\|_{L_{\xi,\tau}^2} \\ &= c\|\prod_{i=1}^4 J^s u_i\|_{L_{x,t}^2} \leq c\prod_{i=1}^4 \|J^s u_i\|_{L_{x,t}^8} \leq c\prod_{i=1}^4 \|u_i\|_{X_{s,b}}, \end{aligned}$$

where in the last step Lemma 2, part ii), with $p = q = 8$ was applied.

Besides $|\xi_{max}| \geq c$ ($\Rightarrow \langle \xi_{max} \rangle \leq c|\xi_{max}|$) we shall assume for the region B that

- i) $|\xi_{min}| \leq 0.99|\xi_{max}|$ or
- ii) $|\xi_{min}| > 0.99|\xi_{max}|$, and there are exactly two indices $i \in \{1, 2, 3, 4\}$ with $\xi_i > 0$.

Then the region B can be splitted again into a finite number of subregions, so that for any of these subregions there exists a permutation π of $\{1, 2, 3, 4\}$ with

$$|\xi| < \xi >^s \prod_{i=1}^4 < \xi_i >^{-s} \leq c |\xi_{\pi(1)} + \xi_{\pi(2)}|^{\frac{1}{2}} |\xi_{\pi(1)} - \xi_{\pi(2)}|^{\frac{1}{2}} < \xi_{\pi(3)} >^{-\frac{3s}{2}} < \xi_{\pi(4)} >^{-\frac{3s}{2}}.$$

Assume $\pi = id$ for the sake of simplicity now. Then we get the upper bound

$$\begin{aligned} & \| < \tau - \xi^3 >^{b'} \int d\nu |\xi_1 + \xi_2|^{\frac{1}{2}} |\xi_1 - \xi_2|^{\frac{1}{2}} < \xi_3 >^{-\frac{3s}{2}} < \xi_4 >^{-\frac{3s}{2}} \prod_{i=1}^4 < \tau_i - \xi_i^3 >^{-b} f_i(\xi_i, \tau_i) \|_{L_{\xi, \tau}^2} \\ & = c \| (I^{\frac{1}{2}} I_-^{\frac{1}{2}} (J^s u_1, J^s u_2)) (J^{-\frac{s}{2}} u_3) (J^{-\frac{s}{2}} u_4) \|_{X_{0, b'}} \end{aligned}$$

To estimate the latter expression, we fix some Sobolev- and Hölderexponents:

- i) $\frac{1}{q_0} = \frac{1}{2} - b'$ so that $L_t^{q_0}(L_x^2) \subset X_{0, b'}$,
- ii) $\frac{2}{p} = \frac{1}{q_0} - \frac{1}{2} = -b'$,
- iii) $\frac{1}{q} = \frac{1}{2} - \frac{2}{p} = \frac{1}{2} + b'$ so that by Lemma 2 $\|J^{\frac{1}{p}} u\|_{L_t^p(L_x^q)} \leq c \|u\|_{X_{0, b'}}$,
- iv) $\epsilon = \frac{1}{p} + \frac{3s}{2} > \frac{1}{q}$ (since $s > \frac{1}{3} + b'$) so that $H_x^{\epsilon, q} \subset L_x^\infty$.

Now we have

$$\begin{aligned} & \| (I^{\frac{1}{2}} I_-^{\frac{1}{2}} (J^s u_1, J^s u_2)) (J^{-\frac{s}{2}} u_3) (J^{-\frac{s}{2}} u_4) \|_{X_{0, b'}} \\ & \leq c \| (I^{\frac{1}{2}} I_-^{\frac{1}{2}} (J^s u_1, J^s u_2)) (J^{-\frac{s}{2}} u_3) (J^{-\frac{s}{2}} u_4) \|_{L_t^{q_0}(L_x^2)} \\ & \leq c \| I^{\frac{1}{2}} I_-^{\frac{1}{2}} (J^s u_1, J^s u_2) \|_{L_{x_t}^2} \| J^{-\frac{s}{2}} u_3 \|_{L_t^p(L_x^\infty)} \| J^{-\frac{s}{2}} u_4 \|_{L_t^p(L_x^\infty)}. \end{aligned}$$

Now by Lemma 1 the first factor can be controlled by $c \|u_1\|_{X_{s, b}} \|u_2\|_{X_{s, b}}$, while for the second we have the upper bound

$$c \| J^{-\frac{3s}{2} + \epsilon} J^s u_3 \|_{L_t^p(L_x^q)} = c \| J^{\frac{1}{p}} J^s u_3 \|_{L_t^p(L_x^q)} \leq c \|u_3\|_{X_{s, b}}.$$

The third factor can be treated in precisely the same way. So for the contributions of the region B we have obtained the desired bound.

Finally we consider the remaining region C : Here the $|\xi_i|$, $1 \leq i \leq 4$, are all very close together and $\geq c < \xi_i >$. Moreover, at least three of the variables ξ_i have the same sign. Thus for the quantity $c.q.$ controlled by the expressions $< \tau - \xi^3 >$, $< \tau_i - \xi_i^3 >$, $1 \leq i \leq 4$, we have in this region:

$$c.q. := |\xi^3 - \sum_{i=1}^4 \xi_i^3| \geq c \sum_{i=1}^4 < \xi_i >^3 \geq c < \xi >^3$$

and hence, since $s > \frac{1}{3} + b'$ is assumed,

$$|\xi| < \xi >^s \prod_{i=1}^4 < \xi_i >^{-s} \leq c (< \tau - \xi^3 >^{-b'} + \sum_{i=1}^4 < \tau_i - \xi_i^3 >^{-b'} \chi_{C_i}),$$

where in the subregion C_i , $1 \leq i \leq 4$, the expression $\langle \tau_i - \xi_i^3 \rangle$ is dominant. The first contribution can be estimated by

$$\begin{aligned} & c \left\| \int d\nu \prod_{i=1}^4 \langle \tau_i - \xi_i^3 \rangle^{-b} f_i(\xi_i, \tau_i) \right\|_{L_{\xi, \tau}^2} \\ &= c \left\| \prod_{i=1}^4 J^s u_i \right\|_{L_{x, t}^2} \leq c \prod_{i=1}^4 \|J^s u_i\|_{L_{x, t}^8} \leq c \prod_{i=1}^4 \|u_i\|_{X_{s, b}}, \end{aligned}$$

where we have used Lemma 2, part ii). For the contribution of the subregion C_1 we take into account that $\langle \tau_1 - \xi_1^3 \rangle = \max\{\langle \tau - \xi^3 \rangle, \langle \tau_i - \xi_i^3 \rangle, 1 \leq i \leq 4\}$, which gives

$$\langle \tau - \xi^3 \rangle^{b+b'} |\xi| \langle \xi \rangle^s \prod_{i=1}^4 \langle \xi_i \rangle^{-s} \leq c \langle \tau_1 - \xi_1^3 \rangle^b.$$

So, for this contribution we get the upper bound

$$\begin{aligned} & c \left\| \langle \tau - \xi^3 \rangle^{-b} \int d\nu \langle \tau_1 - \xi_1^3 \rangle^b \prod_{i=1}^4 \langle \tau_i - \xi_i^3 \rangle^{-b} f_i(\xi_i, \tau_i) \right\|_{L_{\xi, \tau}^2} \\ & \leq c \|\mathcal{F}^{-1} f_1 \prod_{i=2}^4 J^s u_i\|_{X_{0, -b}} \leq c \|\mathcal{F}^{-1} f_1 \prod_{i=2}^4 J^s u_i\|_{L_{x, t}^{\frac{8}{7}}} \\ & \leq c \|\mathcal{F}^{-1} f_1\|_{L_{x, t}^2} \prod_{i=2}^4 \|J^s u_i\|_{L_{x, t}^8} \leq c \prod_{i=1}^4 \|u_i\|_{X_{s, b}}. \end{aligned}$$

Here we have used the dual version of the L^8 -Strichartz estimate, Hölder and the estimate itself. For the remaining subregions C_i the same argument applies. $\square$

Corollary 2 For $s \geq 0$, $-\frac{1}{2} < b' < -\frac{1}{3}$ and $b > \frac{1}{2}$ the estimate (1) holds true.

Proof: For $s = 0$ this is contained in the above theorem, while for $s > 0$ one only has to use $\langle \xi \rangle \leq c \prod_{i=1}^4 \langle \xi_i \rangle$. $\square$

Acknowledgement: I want to thank Professor Hartmut Pecher for numerous helpful conversations.

References

- [B93] Bourgain, J.: Fourier transform restriction phenomena for certain lattice subsets and applications to nonlinear evolution equations, GAFA 3 (1993), 107 - 156 and 209 - 262
- [GTV97] Ginibre, J., Tsutsumi, Y., Velo, G.: On the Cauchy-Problem for the Zakharov-System, J. of Functional Analysis 151 (1997), 384 - 436
- [KPV91] Kenig, C. E., Ponce, G., Vega, L.: Oscillatory Integrals and Regularity of Dispersive Equations, Indiana Univ. Math. J. 40 (1991), 33 - 69
- [KPV93] Kenig, Ponce, Vega: Wellposedness and scattering results for the generalized Korteweg-de Vries equation via the contraction principle, CPAM 46 (1993), 527 - 620
- [KPV96] Kenig, Ponce, Vega: A bilinear estimate with applications to the KdV equation, J. of the AMS 9 (1996), 573 - 603